

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 June 2005 (09.06.2005)

PCT

(10) International Publication Number
WO 2005/051921 A1

(51) International Patent Classification⁷: **C07D 239/42**

(74) Agent: BRAINARD, Charles, R.; Kenyon & Kenyon,
One Broadway, New York, NY 10004-1050 (US).

(21) International Application Number:

PC/T/US2004/039469

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AH, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PI, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:

24 November 2004 (24.11.2004)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/525,128 24 November 2003 (24.11.2003) US
60/534,479 5 January 2004 (05.01.2004) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GI, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except BB, US): TEVA PHARMACEUTICAL INDUSTRIES LTD. [IL/IL]; 5 Basel Street, P.O. Box 3190, 49131 Petah Tiqva (IL).

(71) Applicant (for BB only): TEVA PHARMACEUTICALS USA, INC. [US/US]; 1090 Horsham Road, P.O. Box 1090, North Wales, PA 19454 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): NIDDAM-HILDESHEIM, Valerie [IL/IL]; POB 21, 40696 Ein Vered (IL); ARONHIME, Judith [IL/IL]; Rehov Harav Maor Josef 5a, 76217 Rehovot (IL).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2005/051921 A1

(54) Title: CRYSTALLINE AMMONIUM SALTS OF ROSUVASTATIN

(57) Abstract: Provided are alkyl ammonium crystalline salts of rosuvastatin that provide for purification of rosuvastatin and its pharmaceutically acceptable salts.

CRYSTALLINE AMMONIUM SALTS OF ROSUVASTATIN

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application Nos.

- 5 60/525,128 filed on November 24, 2003 and 60/534,479 filed on January 5, 2004, the disclosures of which are incorporated by reference in their entirety herein.

FIELD OF THE INVENTION

The present invention relates to ammonium salts of rosuvastatin.

10

BACKGROUND OF THE INVENTION

Statins are currently the most therapeutically effective drugs available for reducing low-density lipoprotein (LDL) particle concentration in the blood stream of patients at risk for cardiovascular disease. Thus, statins are used in the treatment of


15 hypercholesterolemia, hyperlipoproteinemia, and atherosclerosis. A high level of LDL in the bloodstream has been linked to the formation of coronary lesions that obstruct the flow of blood and can rupture and promote thrombosis. Goodman and Gilman, The Pharmacological Basis of Therapeutics, page 879 (9th Ed. 1996).

- 20 Statins inhibit cholesterol biosynthesis in humans by competitively inhibiting the 3-hydroxy-3-methyl-glutaryl-coenzyme A ("HMG-CoA") reductase enzyme. HMG-CoA reductase catalyzes the conversion of HMG to mevalonate, which is the rate-determining step in the biosynthesis of cholesterol. Decreased production of cholesterol causes an increase in the number of LDL receptors and corresponding reduction in the concentration of LDL particles in the bloodstream. Reduction in the
- 25 LDL level in the bloodstream reduces the risk of coronary artery disease. J.A.M.A. 1984, 251, 351-74.

- Currently available statins include lovastatin, simvastatin, pravastatin, fluvastatin, cerivastatin and atorvastatin. Lovastatin (disclosed in U.S. Pat. No. 4,231,938) and simvastatin (disclosed in U.S. Pat. No. 4,444,784) are administered in
- 30 the lactone form. After absorption, the lactone ring is opened in the liver by chemical or enzymatic hydrolysis, and the active hydroxy acid is generated. Pravastatin (disclosed in U.S. Pat. No. 4,346,227) is administered as the sodium salt. Fluvastatin (disclosed in U.S. Pat. No. 4,739,073) and cerivastatin (disclosed in U.S.

Pat. No. 5,006,530 and 5,177,080), also administered as the sodium salt, are entirely synthetic compounds that are in part structurally distinct from the fungal derivatives of this class that contain a hexahydronaphthalene ring. Atorvastatin and two new "superstatins," rosuvastatin and pitavastatin, are administered as calcium salts.

- 5 Rosuvastatin calcium (7-[4-(4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methylsulfonylamino)pyrimidin-5-yl]-(3R,5S)-dihydroxy-(E)-6-heptenoic acid) is an HMG-CoA reductase inhibitor, developed by Shionogi for the once daily oral treatment of hyperlipidaemia (Ann Rep, Shionogi, 1996; Direct communications, Shionogi, 8 Feb 1999 & 25 Feb 2000). Rosuvastatin calcium is a so-called
- 10 superstatin, which can lower LDL-cholesterol and triglycerides more effectively than first generation drugs. Rosuvastatin calcium has the following chemical formula:



- 15 Rosuvastatin calcium is marketed under the name CRESTOR for treatment of a mammal such as a human. According to the maker of CRESTOR, it is administered in a daily dose of from about 5mg to about 40 mg. For patients requiring less aggressive LDL-C reductions or who have pre-disposing factors for myopathy, the 5mg dose is recommended, while 10 mg dose is recommended for the average patient,
- 20 20 mg dose for patients with marked hyper-cholesterolemia and aggressive lipid targets (>190 mg/dL), and the 40 mg dose for patients who have not been responsive to lower doses.

- U.S. patent No. 5,260,440 discloses and claims rosuvastatin, its calcium salt (2:1), and its lactone form. The process of the '440 patent prepares rosuvastatin by
- 25 reacting 4-(4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methylsulfonylamino)-5-pyrimidinecarbaldehyde with methyl (3R)-3-(tert-butyldimethylsilyloxy)-5-oxo-6-triphenylphosphoranylidene hexanate in acetonitrile under reflux. The silyl group is

then cleaved with hydrogen fluoride, followed by reduction with NaBH_4 to obtain a methyl ester of rosuvastatin.

The ester is then hydrolyzed with sodium hydroxide in ethanol at room temperature, followed by removal of ethanol and addition of ether, to obtain the sodium salt of rosuvastatin. The sodium salt is then converted to the calcium salt with a multi-step process. The sodium salt is dissolved in water and maintained under a nitrogen atmosphere. Calcium chloride is then added to the solution, resulting in precipitation of rosuvastatin calcium (2:1). The process for preparation of the intermediates disclosed in the '440 patent is incorporated herein by reference.

U.S. Pat. No. 6,316,460 discloses a pharmaceutical composition of rosuvastatin. The pharmaceutical compositions contain rosuvastatin or its salt and a multivalent tribasic phosphate salt.

WO 01/60804 discloses various ammonium, lithium and magnesium salts of rosuvastatin.

Even though ammonium salts of rosuvastatin are not likely to be used for administration to a patient, they provide a method for purifying rosuvastatin through crystallization. There is a need in the art for additional salts of rosuvastatin that allow for purification of rosuvastatin through crystallization, particularly since rosuvastatin calcium as marketed under CRESTOR is highly amorphous.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides for crystalline rosuvastatin isopropylammonium salt.

In another aspect, the present invention provides for a crystalline form of rosuvastatin isopropylammonium salt having an X-ray powder diffraction pattern with peaks at 6.2, 18.8, 19.3, 20.6 and 22.3 ± 0.2 deg. 2-theta.

In another aspect, the present invention provides for a process for preparing the above crystalline form of rosuvastatin comprising combining a solution of rosuvastatin acid in an organic solvent with isopropylamine to precipitate the crystalline form and recovering the crystalline form.

In another aspect, the present invention provides for a process for preparing the above crystalline form comprising combining a solution of rosuvastatin acid in acetonitrile or ethyl acetate with isopropylamine followed by stirring at a temperature

of about 30°C, followed by cooling to a temperature of less than about 10°C to precipitate the crystalline form and recovering the crystalline form.

In another aspect, the present invention provides for a process for purifying rosuvastatin calcium comprising

- 5 a) converting rosuvastatin calcium salt to rosuvastatin acid;
 b) converting rosuvastatin acid to the above isopropylammonium salt;
 c) converting the isopropylammonium salt to rosuvastatin calcium.

In another aspect, the present invention provides for a process for purifying rosuvastatin calcium comprising

- 10 a) combining rosuvastatin calcium with water to obtain a mixture;
 b) combining the mixture with an acid to obtain a solution of rosuvastatin acid;
 c) extracting the solution with a water immiscible solvent to obtain an organic phase;
15 d) evaporating the organic phase to obtain a residue;
 e) combining the residue with an organic solvent and isopropylamine to obtain the above isopropyl ammonium salt;
 f) cooling the organic solvent to precipitate the isopropyl ammonium salt; and
 g) converting the isopropyl ammonium salt to the calcium salt.

- 20 In another aspect, the present invention provides for a crystalline rosuvastatin cyclohexylammonium salt.

In another aspect, the present invention provides for a crystalline form of rosuvastatin cyclohexylammonium salt having an X-ray powder diffraction pattern characterized by peaks at 8.7, 9.5, 15.3, 19.6 and 20.8 ± 0.2 deg. 2-theta.

- 25 In another aspect, the present invention provides for a process for preparing the crystalline form of rosuvastatin cyclohexylammonium salt comprising combining a solution of rosuvastatin acid in an organic solvent with cyclohexylamine to precipitate the crystalline form and recovering the crystalline form.

- In another aspect, the present invention provides for a process for preparing
30 the above crystalline form comprising combining a solution of rosuvastatin acid in ethyl acetate with cyclohexylamine followed by stirring at a temperature of about 30°C, followed by cooling to a temperature of less than about 10°C to precipitate the crystalline form and recovering the crystalline form.

In another aspect, the present invention provides for a process for purifying rosuvastatin calcium comprising

- a) converting rosuvastatin calcium salt to rosuvastatin acid;
- b) converting rosuvastatin acid to the above cyclohexylammonium salt;
- 5 c) converting the isopropylammonium salt to rosuvastatin calcium.

In another aspect, the present invention provides for a process for purifying rosuvastatin calcium comprising

- a) combining rosuvastatin calcium with water to obtain a mixture;
- b) combining the mixture with an acid to obtain a solution of rosuvastatin
- 10 acid;
- c) extracting the solution with a water immiscible solvent to obtain an organic phase;
- d) evaporating the organic phase to obtain a residue;
- e) combining the residue with an organic solvent and the cyclohexylamine to
- 15 obtain the above cyclohexylamine salt;
- f) cooling the organic solvent to precipitate the cyclohexylammonium salt; and
- g) converting the cyclohexylammonium salt to the calcium salt.

In another aspect, the present invention provides for a process for purifying rosuvastatin calcium comprising

- 20 a) combining rosuvastatin calcium with water to obtain a mixture;
- b) combining the mixture with an acid to obtain a solution of rosuvastatin acid;
- c) extracting the solution with a water immiscible solvent to obtain an organic phase;
- 25 d) evaporating the organic phase to obtain a residue;
- e) combining the residue with an organic solvent and an amine to obtain the ammonium salt;
- f) cooling the organic solvent to precipitate the ammonium salt; and
- g) converting the ammonium salt to the calcium salt.

30

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is an X-ray Powder Diffraction pattern of rosuvastatin isopropylammonium Form A1.

Figure 2 is an X-ray powder diffraction pattern of rosuvastatin cyclohexylammonium Form B.

5 DETAILED DESCRIPTION OF THE INVENTION

One approach for formulation is to use a crystalline form instead of amorphous form in order to minimize problems generally associated with amorphous form. Another approach for formulation is to purify amorphous form, so that the greater purity imparts additional stability to amorphous form.

10 An approach to purifying an amorphous active pharmaceutical ingredient (“API”) is using a crystalline salt as an intermediate. Crystallization often allows for purification of an API since impurities may be left in the solution after crystallization, or be washed away as insoluble impurities before crystallization. Additional impurities may also be removed by washing the crystals.

15 The present invention provides crystalline isopropylammonium and cyclohexylammonium salts of rosuvastatin and procedures for their preparation. These salts are useful intermediates for purification of the final product, which is often administered as a calcium salt. The ammonium salt intermediate is particularly useful since it is crystalline, and not highly amorphous like rosuvastatin calcium. For
20 example, the maker of CRESTOR refers to CRESTOR as a white amorphous powder. Amorphous form of an active pharmaceutical ingredient often exhibit lower stability and greater reactivity. Amorphous material may contain impurities, and crystallization as a salt, such as an ammonium salt, allows for purifying the amorphous material.

The purification can start with the calcium salt, followed by its conversion to the acid, which is then converted to the ammonium salt. The ammonium salt may then be converted to a highly pure calcium salt. The purification can also start from other salt or acid forms of rosuvastatin, followed by conversion to the ammonium salt, followed by conversion to the calcium salt. *See e.g.* WO 04/014872.

The XRD peaks of the two crystalline forms are as follows:

30 Typical XRD peaks - deg. 2-theta ± 0.2 (The main characteristic peaks are underlined)

<u>Form A1</u>	<u>Form B</u>
6.0	<u>8.7</u>
<u>6.2</u>	<u>9.5</u>

10.2	10.7
10.6	14.3
13.4	14.7
15.6	<u>15.3</u>
16.6	17.4
18.2	18.0
<u>18.8</u>	<u>19.6</u>
<u>19.3</u>	<u>20.8</u>
<u>20.6</u>	22.1
<u>22.3</u>	28.0
27.3	
28.8	

The present invention provides crystalline rosuvastatin-isopropylammonium salt.

Crystalline rosuvastatin isopropylammonium salt of the present invention
 5 (Form A1) has an X-ray powder diffraction pattern (Figure 1) with principal characteristic peaks at 6.2, 18.8, 19.3, 20.6 and 22.3 ± 0.2 deg. 2-theta. Other characteristic peaks are at 6.0, 10.2, 10.6, 13.4, 15.6, 16.6, 18.2, 27.3 and 28.8 ± 0.2 deg. 2-theta. This Form was previously referred to as Form A in the priority application of the present invention.

10 Form A1 is generally prepared by combining a solution of rosuvastatin acid in an organic solvent with isopropylamine to precipitate the crystalline form and recovering the crystalline form. After combining, the reaction mixture is preferably stirred at a temperature of about 20°C to about 40°C, more preferably at a temperature of about 30°C for a few hours. If precipitation does not occur, the solution is
 15 preferably cooled to a temperature of less than about 10°C. Preferably, the solvent is acetonitrile or ethyl acetate.

The present invention provides crystalline form of rosuvastatin cyclohexylammonium salt.

Crystalline form of rosuvastatin cyclohexylammonium salt of the present
 20 invention (Form B) has an XRPD pattern (Figure 2) with principal peaks at 8.7, 9.5,

15.3, 19.6 and 20.8 ± 0.2 deg. 2-theta. Other peaks are at 10.7, 14.3, 14.7, 17.4, 18.0, 22.1 and 28.1 ± 0.2 deg. 2-theta.

Form B is generally prepared by combining a solution of rosuvastatin acid in a suitable organic solvent with cyclohexylamine to precipitate the crystalline form and recovering the crystalline form. After combining, the reaction mixture is preferably stirred at a temperature of about 20°C to about 40°C, more preferably at a temperature of about 30°C for a few hours. If precipitation does not occur, the solution is preferably cooled to a temperature of less than about 10°C. Preferably, the organic solvent is ethyl acetate.

10 Form B may be prepared from rosuvastatin calcium as described above in regard to Form A1.

The present invention encompasses embodiments where the amine, rosuvastatin acid/calcium and the organic solvent are added simultaneously, though it is preferred to first obtain a solution of the acid/calcium salt in the solvent.

15 The above crystalline forms may be recovered by conventional techniques such as filtration, decanting and centrifugation. The resulting salts may optionally be dried to remove residual solvent, such as by drying under reduced pressure (below 1 atmosphere or preferably below about 100mmHg).

The starting material used in the present invention to obtain the salt is 20 rosuvastatin acid. The acid can be obtained from rosuvastatin calcium. The acid may be isolated and subsequently used, or be used in a one pot procedure where the calcium salt is converted to the amine without isolation of the acid.

In one embodiment, rosuvastatin calcium of amorphous nature is purified by conversion to the acid, which is converted to the above ammonium salts, which is 25 then converted back to amorphous rosuvastatin calcium. The rosuvastatin calcium obtained at the end has a higher purity than the rosuvastatin calcium used as starting material. This scheme can also be used with other pharmaceutically acceptable salts of rosuvastatin which are also amorphous.

In one embodiment, rosuvastatin calcium is combined with water or with a 30 mixture of water and a water-miscible solvent. A preferred mixture is about a 1:2 to about a 2:1 mixture of water and acetonitrile, more preferably a 1:1 mixture (by volume). The weight to volume ratio of the calcium salt is preferably of about 1:10 to about 1:40 Kg/L. The ratio might vary depending on the solvent used, but is of such

that a solution would be formed after formation of the acid, and of sufficient concentration to allow for optimal extraction. The acid form may be obtained from the calcium salt by addition of an acid, such as HCl, HBr, or sulfuric acid. After addition of the acid, the reaction mixture may be stirred.

5 The resulting solution is then extracted with a water immiscible solvent. Preferably, the water immiscible solvent is ethyl acetate. The rosuvastatin acid moves into the organic phase. After extraction, the organic phase may be concentrated, preferably by evaporation at a pressure below about 760mmHg, more preferably below about 100mmHg. The temperature can be increased to accelerate the
10 evaporation process. In this embodiment, the resulting residue is then added to an organic solvent, followed by addition of the amine to precipitate the ammonium salt. After the addition, the reaction mixture is preferably stirred at a temperature of about 20°C to about 40°C, more preferably at a temperature of about 30°C for a few hours. Preferably, the organic solvent is ethyl acetate. Preferably, the amine is
15 isopropylamine or cyclohexylamine. When the amine is isopropylamine, the ammonium salt is crystalline rosuvastatin isopropylammonium salt form A1. When the amine is cyclohexylamine, the ammonium salt is crystalline rosuvastatin cyclohexylamine salt form B. Precipitation is preferably induced by cooling as described above.

20 After obtaining the above ammonium salts, the ammonium salts can be converted to rosuvastatin calcium for example by reaction with an aqueous basic solution of sodium or potassium hydroxide (about 8%), followed by reaction with a source of calcium such as calcium chloride or calcium oxide. The calcium salt may be recovered in the solid state by removing the solvent under reduced pressure or by
25 precipitation.

 Many processes of the present invention involve crystallization/precipitation out of a particular solvent, i.e., obtaining a solid material from a solution. One skilled in the art would appreciate that the conditions concerning crystallization may be modified without affecting the form of the polymorph obtained. For example, when
30 mixing rosuvastatin in a solvent to form a solution, warming of the mixture may be necessary to completely dissolve the starting material. If warming does not clarify the mixture, the mixture may be diluted or filtered. To filter, the hot mixture may be passed through paper, glass fiber or other membrane material, or a clarifying agent such as celite. Depending upon the equipment used and the concentration and

temperature of the solution, the filtration apparatus may need to be preheated to avoid premature crystallization.

The conditions may also be changed to induce precipitation. A preferred way of inducing precipitation is to reduce the solubility of the solvent. The solubility of the solvent may be reduced, for example, by cooling the solvent.

In one embodiment, an anti-solvent is added to a solution to decrease its solubility for a particular compound, thus resulting in precipitation. Another way of accelerating crystallization is by seeding with a crystal of the product or scratching the inner surface of the crystallization vessel with a glass rod. Other times, crystallization may occur spontaneously without any inducement. The present invention encompasses both embodiments where crystallization of a particular form of rosuvastatin occurs spontaneously or is induced/accelerated, unless if such inducement is critical for obtaining a particular form.

Pharmaceutical compositions can be prepared as medicaments to be administered orally, parenterally, rectally, transdermally, buccally, or nasally. Suitable forms for oral administration include tablets, compressed or coated pills, dragees, sachets, hard or gelatin capsules, sub-lingual tablets, syrups and suspensions. Suitable forms of parenteral administration include an aqueous or non-aqueous solution or emulsion, while for rectal administration suitable forms for administration include suppositories with hydrophilic or hydrophobic vehicle. For topical administration the invention provides suitable transdermal delivery systems known in the art, and for nasal delivery there are provided suitable aerosol delivery systems known in the art.

Pharmaceutical formulations of the present invention contain a pharmaceutically acceptable salt of rosuvastatin in the solid state. In addition to the active ingredient(s), the pharmaceutical compositions of the present invention may contain one or more excipients or adjuvants. Selection of excipients and the amounts to use may be readily determined by the formulation scientist based upon experience and consideration of standard procedures and reference works in the field.

Diluents increase the bulk of a solid pharmaceutical composition, and may make a pharmaceutical dosage form containing the composition easier for the patient and care giver to handle. Diluents for solid compositions include, for example, microcrystalline cellulose (e.g. Avicel[®]), microfine cellulose, lactose, starch, pregelatinized starch, calcium carbonate, calcium sulfate, sugar, dextrans, dextrin, dextrose, dibasic calcium phosphate dihydrate, tribasic calcium phosphate, kaolin,

magnesium carbonate, magnesium oxide, maltodextrin, mannitol, polymethacrylates (e.g. Eudragit®), potassium chloride, powdered cellulose, sodium chloride, sorbitol and talc.

5 Solid pharmaceutical compositions that are compacted into a dosage form, such as a tablet, may include excipients whose functions include helping to bind the active ingredient and other excipients together after compression. Binders for solid pharmaceutical compositions include acacia, alginic acid, carbomer (e.g. carbopol), carboxymethylcellulose sodium, dextrin, ethyl cellulose, gelatin, guar gum, hydrogenated vegetable oil, hydroxyethyl cellulose, hydroxypropyl cellulose (e.g. Klucel®), hydroxypropyl methyl cellulose (e.g. Methocel®), liquid glucose, 10 magnesium aluminum silicate, maltodextrin, methylcellulose, polymethacrylates, povidone (e.g. Kollidon®, Plasdone®), pregelatinized starch, sodium alginate and starch.

The dissolution rate of a compacted solid pharmaceutical composition in the 15 patient's stomach may be increased by the addition of a disintegrant to the composition. Disintegrants include alginic acid, carboxymethylcellulose calcium, carboxymethylcellulose sodium (e.g. Ac-Di-Sol®, Primellose®), colloidal silicon dioxide, croscarmellose sodium, crospovidone (e.g. Kollidon®, Polyplasdone®), guar gum, magnesium aluminum silicate, methyl cellulose, microcrystalline cellulose, 20 polacrillin potassium, powdered cellulose, pregelatinized starch, sodium alginate, sodium starch glycolate (e.g. Explotab®) and starch.

Glidants can be added to improve the flowability of a non-compacted solid composition and to improve the accuracy of dosing. Excipients that may function as glidants include colloidal silicon dioxide, magnesium trisilicate, powdered cellulose, 25 starch, talc and tribasic calcium phosphate.

When a dosage form such as a tablet is made by the compaction of a powdered composition, the composition is subjected to pressure from a punch and dye. Some excipients and active ingredients have a tendency to adhere to the surfaces of the punch and dye, which can cause the product to have pitting and other surface 30 irregularities. A lubricant can be added to the composition to reduce adhesion and ease the release of the product from the dye. Lubricants include magnesium stearate, calcium stearate, glyceryl monostearate, glyceryl palmitostearate, hydrogenated castor oil, hydrogenated vegetable oil, mineral oil, polyethylene glycol, sodium benzoate, sodium lauryl sulfate, sodium stearyl fumarate, stearic acid, talc and zinc stearate.

Flavoring agents and flavor enhancers make the dosage form more palatable to the patient. Common flavoring agents and flavor enhancers for pharmaceutical products that may be included in the composition of the present invention include maltol, vanillin, ethyl vanillin, menthol, citric acid, fumaric acid, ethyl maltol, and tartaric acid.

Solid and liquid compositions may also be dyed using any pharmaceutically acceptable colorant to improve their appearance and/or facilitate patient identification of the product and unit dosage level.

In liquid pharmaceutical compositions of the present invention, nateglinide and any other solid excipients are dissolved or suspended in a liquid carrier such as water, vegetable oil, alcohol, polyethylene glycol, propylene glycol or glycerin. Liquid pharmaceutical compositions may contain emulsifying agents to disperse uniformly throughout the composition an active ingredient or other excipient that is not soluble in the liquid carrier. Emulsifying agents that may be useful in liquid compositions of the present invention include, for example, gelatin, egg yolk, casein, cholesterol, acacia, tragacanth, chondrus, pectin, methyl cellulose, carbomer, cetostearyl alcohol and cetyl alcohol.

Liquid pharmaceutical compositions of the present invention may also contain a viscosity enhancing agent to improve the mouth-feel of the product and/or coat the lining of the gastrointestinal tract. Such agents include acacia, alginic acid bentonite, carbomer, carboxymethylcellulose calcium or sodium, cetostearyl alcohol, methyl cellulose, ethylcellulose, gelatin guar gum, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, maltodextrin, polyvinyl alcohol, povidone, propylene carbonate, propylene glycol alginate, sodium alginate, sodium starch glycolate, starch tragacanth and xanthan gum.

Sweetening agents such as sorbitol, saccharin, sodium saccharin, sucrose, aspartame, fructose, mannitol and invert sugar may be added to improve the taste. Preservatives and chelating agents such as alcohol, sodium benzoate, butylated hydroxy toluene, butylated hydroxyanisole and ethylenediamine tetraacetic acid may be added at levels safe for ingestion to improve storage stability.

According to the present invention, a liquid composition may also contain a buffer such as gluconic acid, lactic acid, citric acid or acetic acid, sodium gluconate, sodium lactate, sodium citrate or sodium acetate.

Selection of excipients and the amounts used may be readily determined by the formulation scientist based upon experience and consideration of standard procedures and reference works in the field.

The solid compositions of the present invention include powders, granulates, aggregates and compacted compositions. The dosages include dosages suitable for oral, buccal, rectal, parenteral (including subcutaneous, intramuscular, and intravenous), inhalant and ophthalmic administration. Although the most suitable administration in any given case will depend on the nature and severity of the condition being treated, the most preferred route of the present invention is oral. The dosages may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the pharmaceutical arts.

Dosage forms include solid dosage forms like tablets, powders, capsules, suppositories, sachets, troches and lozenges, as well as liquid syrups, suspensions and elixirs.

The dosage form of the present invention may be a capsule containing the composition, preferably a powdered or granulated solid composition of the invention, within either a hard or soft shell. The shell may be made from gelatin and optionally contain a plasticizer such as glycerin and sorbitol, and an opacifying agent or colorant.

The active ingredient and excipients may be formulated into compositions and dosage forms according to methods known in the art.

A composition for tableting or capsule filling may be prepared by wet granulation. In wet granulation, some or all of the active ingredients and excipients in powder form are blended and then further mixed in the presence of a liquid, typically water, that causes the powders to clump into granules. The granulate is screened and/or milled, dried and then screened and/or milled to the desired particle size. The granulate may then be tableted, or other excipients may be added prior to tableting, such as a glidant and/or a lubricant.

A tableting composition may be prepared conventionally by dry blending. For example, the blended composition of the actives and excipients may be compacted into a slug or a sheet and then comminuted into compacted granules. The compacted granules may subsequently be compressed into a tablet.

As an alternative to dry granulation, a blended composition may be compressed directly into a compacted dosage form using direct compression techniques. Direct

compression produces a more uniform tablet without granules. Excipients that are particularly well suited for direct compression tableting include microcrystalline cellulose, spray dried lactose, dicalcium phosphate dihydrate and colloidal silica. The proper use of these and other excipients in direct compression tableting is known to those in the art with experience and skill in particular formulation challenges of direct compression tableting.

A capsule filling of the present invention may comprise any of the aforementioned blends and granulates that were described with reference to tableting, however, they are not subjected to a final tableting step.

Preferred dosage is from about 5mg to about 80mg per day, more preferably about 5mg to about 40mg per day, with 5mg, 10mg, 20mg, 40mg and 80 mg tablets for once a day being preferred. These tablets may have the following inactive ingredients: microcrystalline cellulose NF, lactose monohydrate NF, tribasic calcium phosphate NF, croscopovidone NF, magnesium stearate NF, hypromellose NF, triacetin NF and titanium dioxide USP.

Also provided are formulations of highly pure calcium salt of rosuvastatin and its methods of administration.

X-Ray powder diffraction data were obtained by conventional methods using a SCINTAG powder X-Ray diffractometer model XTRA equipped with a solid state detector. Copper radiation of 1.5418 lambda was used. A round aluminum sample holder with zero background was used. All peak positions are within 0.2 degrees two theta.

Procedures

Example 1: Preparation of Rosuvastatin iso-Propylammonium salt in ACN:

A 100 ml three necked flask equipped with a mechanical stirrer was charged with rosuvastatin Ca salt (3g), water (30 ml) and acetonitrile (30 ml).

6 ml of HCl 1N was added portion-wise at room temperature. The mixture was stirred at room temperature for 1 hour. The clear solution was then extracted with 3 portions of 20 ml ethyl acetate. The organic phase was concentrated under reduced pressure of 10-100 mm Hg at 45°C to obtain a residue (4 g), which contains rosuvastatin acid.

Acetonitrile (30 mL) was added to the residue, the solution was stirred at room temperature for 5 minutes. 0.36 g of iso-propylamine were added dropwise and the solution was stirred at 30°C for 90 minutes, then cooled at 0°-2°C (ice-water bath), and stirred for 2 hours. The iso-propylammonium salt was collected by filtration
5 under reduced pressure to obtain white crystals. (1.6g , 49.4%) in crystal form A1.

Example 2: Preparation of Rosuvastatin iso-Propylammonium salt in Ethyl acetate:

A 100 ml three necked flask equipped with a mechanical stirrer was charged with rosuvastatin Ca salt (3g), water (30 ml) and acetonitrile (30 ml).
10 6 ml of HCl 1N was added portion-wise at room temperature. The mixture was stirred at room temperature for 1 hour. The clear solution was then extracted with 3 portions of 20ml ethyl acetate. 6g of anhydrous Na₂SO₄ was added to the organic phase, the mixture was then stirred for 5 minutes, filtered and concentrated under reduced pressure of 10-100 mm Hg at 45°C to obtain a residue (4 g) which contains
15 rosuvastatin acid.
15 ml of Ethyl acetate was added to the residue, the solution was stirred at room temperature for 5 minutes. 0.36 g of iso-propylamine were added dropwise and the solution was stirred at 30°C for 90 minutes, then cooled at 0°-2°C (ice-water bath) and stirred for 2 hours. The iso-propylammonium salt was collected by filtration under
20 reduced pressure and washed with 10 ml of ethyl acetate to obtain white crystals. (2.0g, 61.7%) in crystal form A1.

Example 3: Same as example 2 except the preparation of the rosuvastatin acid:

A 100 ml three necked flask equipped with a mechanical stirrer was charged with
25 rosuvastatin ca salt (3g) and water (30 ml). 6 ml of HCl 1N was added portion-wise at room temperature. The mixture was stirred at room temperature for 1 hour. 20 ml of Ethyl acetate was added while stirring for 10 minutes to obtain a clear solution. The phases were separated, the aqueous phase was then extracted with 2 portions of 20 ml ethyl acetate. 6g of anhydrous Na₂SO₄ was added to the organic phase, the
30 mixture was then stirred for 5 minutes, filtered and concentrated under reduced pressure of 10-100 mm Hg at 45°C to obtain a residue (4 g) which contains rosuvastatin acid.
15 ml of ethyl acetate was added to the residue, the solution was stirred at room temperature for 5 minutes. 0.36 g of iso-propylamine were added dropwise and the

solution was stirred at 30°C for 90 minutes, then cooled at 0°-2°C (ice-water bath) and stirred for 2 hours. The iso-propylammonium salt was collected by filtration under reduced pressure and washed with 10 ml of ethyl acetate to obtain white crystals (2.5g, 77.2%) in crystal form A1.

5

Example 4: Preparation of Rosuvastatin cyclohexylammonium salt(ROSU-CHA) in Ethyl acetate:

A 100 ml three necked flask equipped with a mechanical stirrer was charged with rosuvastatin Ca salt (3g), water (30 ml) and acetonitrile (30 ml). 6 ml of HCl 1N was
10 added portion-wise at room temperature. The mixture was stirred at room temperature for 1 hour. The clear solution was then extracted with 3 portions of 20 ml ethyl acetate. 6g of anhydrous Na₂SO₄ was added to the organic phase, the mixture was then stirred for 5 minutes, filtered and concentrated under reduced pressure of 10-100 mm Hg at 45°C to obtain a residue (4 g) which contains
15 rosuvastatin acid. 15 ml of ethyl acetate was added to the residue, the solution was stirred at room temperature for 5 minutes. 0.75 ml of cyclohexylamine were added dropwise and the solution was stirred at 30°C for 90 minutes, then cooled at 0°-2°C (ice-water bath) and stirred for 2 hours. The Rosu-CHA salt was collected by filtration under reduced pressure and washed with 10 ml of Ethyl acetate to obtain
20 white crystals (2.9g , 83.5%) in crystal form B.

Example 5-Purification of rosuvastatin calcium by using iso-Propylammonium salt as an intermediate

A 100 ml three necked flask equipped with a mechanical stirrer was charged with
25 Rosuvastatin Ca salt (3g) and water (30 ml).HCl 1N (6 ml) was added portion-wise at room temperature. The mixture was stirred at room temperature for 1 hour. Ethyl acetate (20 ml) was added, while stirring for 10 minutes to obtain a clear solution. The phases were separated. The aqueous phase was then extracted with 2 portions of ethyl acetate (20ml). Anhydrous Na₂SO₄ (6g) was added to the organic phase. The
30 mixture was then stirred for 5 minutes, filtered and concentrated under reduced pressure at 45°C to obtain a residue (4g) which contained Rosuvastatin acid. Ethyl acetate (15 ml) was added to the residue, the solution was stirred at room temperature for 5 minutes. Iso-Propylamine (0.36g) was added dropwise to the solution, and the solution was stirred at 30°C for 90 minutes. The resulting mixture

was cooled at 0°-2°C (ice-water bath) and stirred for 2 hours. The iso-Propylammonium salt was collected by filtration under reduced pressure and washed with ethyl acetate (10 ml) to obtain white crystals (2.5 g, 77.2%) in crystal form A1. A 100 ml three necked flask equipped with a mechanical stirrer was charged at ambient temperature with rosuvastatin iso-Propylammonium salt (2.2 grams) generally prepared according to the above procedure. Water (30 ml), ethanol (22 ml) and sodium hydroxide (0.2 g). The reaction mixture was stirred for 2 hours and the clear solution was filtered under reduced pressure. The mixture was concentrated under reduced pressure at 50°C to obtain a residue. Then water (22 ml) was added and the aqueous phase was washed with EtOAc (2x10 ml). Traces of EtOAc in the aqueous phase were distilled off under reduced pressure at 50°C. Make-up of water was done (10 ml) at the end of the evaporation. To this solution CaCl₂ (0.45g) was added. The reaction mixture was then stirred at ambient temperature for 2 hours, filtered and washed to get 1.5 g (73.6%) of Rosuvastatin calcium.

Having thus described the invention with reference to particular preferred embodiments and illustrated it with Examples, those in the art can appreciate modifications to the invention as described and illustrated that do not depart from the spirit and scope of the invention as disclosed in the specification. The Examples are set forth to aid in understanding the invention but are not intended to, and should not be construed to, limit its scope in any way. The examples do not include detailed descriptions of conventional methods.

What is claimed is:

1. Crystalline rosuvastatin isopropylammonium salt.
- 5 2. A crystalline form of rosuvastatin isopropylammonium salt having an X-ray powder diffraction pattern with peaks at 6.2, 18.8, 19.3, 20.6 and 22.3 ± 0.2 deg. 2-theta.
3. The crystalline form of claim 2, further having peaks at 6.0, 10.2, 10.6, 13.4, 15.6, 16.6, 18.2, 27.3 and 28.8 ± 0.2 deg. 2-theta.
- 10 4. The crystalline form of claim 3, wherein the crystalline form has an X-ray powder diffraction pattern substantially as depicted in Figure 1.
5. A process for preparing a crystalline form of claim 2 comprising combining a solution of rosuvastatin acid in an organic solvent with isopropylamine to precipitate the crystalline form and recovering the crystalline form.
- 15 6. The process of claim 5, wherein precipitation is induced by cooling the solution to a temperature of less than about 10°C.
7. The process of claim 5, wherein the combining is at a temperature of about 20°C to about 40°C.
8. The process of claim 7, wherein the temperature is about 30°C.
- 20 9. The process of claim 5, wherein the solvent is acetonitrile or ethyl acetate.
10. A process for preparing a crystalline form of claim 2 comprising combining a solution of rosuvastatin acid in acetonitrile or ethyl acetate with isopropylamine followed by stirring at a temperature of about 30°C, followed by cooling to a temperature of less than about 10°C to precipitate the
- 25 crystalline form and recovering the crystalline form.
11. A process for purifying rosuvastatin calcium comprising
 - a) converting rosuvastatin calcium salt to rosuvastatin acid;
 - b) converting rosuvastatin acid to the isopropylammonium salt of claim 2;
 - c) converting the isopropylammonium salt to rosuvastatin calcium.
- 30 12. A process for purifying rosuvastatin calcium comprising converting the isopropylammonium salt of claim 2 to rosuvastatin calcium.

13. The process of either of claims 11 or 12, wherein the calcium salt is purified by combining the isopropylammonium salt with an aqueous base to obtain a solution of a sodium or potassium salt, evaporating the solution to obtain a residue, combining the residue with water and a source of calcium to precipitate the calcium salt.
14. A process for purifying rosuvastatin calcium comprising
- a) combining rosuvastatin calcium with water to obtain a mixture;
 - b) combining the mixture with an acid to obtain a solution of rosuvastatin acid;
 - c) extracting the solution with a water immiscible solvent to obtain an organic phase;
 - d) evaporating the organic phase to obtain a residue;
 - e) combining the residue with an organic solvent and isopropylamine to obtain a crystalline form of rosuvastatin isopropylammonium salt having an X-ray powder diffraction pattern with peaks at 6.2, 18.8, 19.3, 20.6 and 22.3 ± 0.2 deg. 2-theta.
 - f) cooling the organic solvent to precipitate the isopropyl ammonium salt; and
 - g) converting the isopropyl ammonium salt to the calcium salt.
15. The process of claim 14, wherein the mixture obtained in step a) further includes a water miscible solvent.
16. The process of claim 15, wherein the water miscible solvent is acetonitrile.
17. The process of claim 16, wherein the ratio between the water and acetonitrile is about 1:2 to about 2:1 by volume.
18. The process of claim 17, wherein the ratio between the water and acetonitrile is about 1:1.
19. The process of claim 14, wherein the weight to volume ratio of the calcium salt is about 1:10 to about 1:40 Kg/L.
20. The process of claim 14, wherein the acid is selected from the group consisting of: hydrochloric, hydrosulfuric, and hydrobromic.
21. The process of claim 14, wherein the water immiscible solvent in step c) is ethyl acetate.

22. The process of claim 14, wherein the evaporation is at a pressure below about 760mmHg.
23. Crystalline rosuvastatin cyclohexylammonium salt.
24. A crystalline form of rosuvastatin cyclohexylammonium salt having an X-ray powder diffraction pattern characterized by peaks at 8.7, 9.5, 15.3, 19.6 and 20.8 \pm 0.2 deg. 2-theta.
25. The crystalline form of claim 24 further having peaks at 10.7, 14.3, 14.7, 17.4, 18.0, 22.1 and 28.1 \pm 0.2 deg. 2-theta.
26. The crystalline form of claim 25 having an X-ray powder diffraction pattern substantially as depicted in Figure 2.
27. A process for preparing a crystalline form of claim 24 comprising combining a solution of rosuvastatin acid in an organic solvent with cyclohexylamine to precipitate the crystalline form and recovering the crystalline form.
28. The process of claim 27, wherein precipitation is induced by cooling the solution to a temperature of less than about 10°C.
29. The process of claim 27, wherein the combining is carried out at a temperature of about 20°C to about 40°C.
30. The process of claim 28, wherein the temperature is about 30°C.
31. The process of claim 27, wherein the organic solvent is ethyl acetate.
32. A process for preparing a crystalline form of claim 24 comprising combining a solution of rosuvastatin acid in ethyl acetate with cyclohexylamine followed by stirring at a temperature of about 30°C, followed by cooling to a temperature of less than about 10°C to precipitate the crystalline form and recovering the crystalline form.
33. A process for purifying rosuvastatin calcium comprising
- a) converting rosuvastatin calcium salt to rosuvastatin acid;
 - b) converting rosuvastatin acid to the cyclohexylammonium salt of claim 24;
 - c) converting the cyclohexylammonium salt to rosuvastatin calcium.
34. A process for purifying rosuvastatin calcium comprising converting the cyclohexylammonium salt of claim 24 to rosuvastatin calcium.
35. The process of either of claims 33 or 34, wherein the calcium salt is prepared by combining the cyclohexylammonium salt with an aqueous base to obtain a solution of a sodium or potassium salt, evaporating the solution to obtain a

- residue, combining the residue with water and a source of calcium to precipitate the calcium salt.
36. A process for purifying rosuvastatin calcium comprising
- 5 a) combining rosuvastatin calcium with water to obtain a mixture;
- b) combining the mixture with an acid to obtain a solution of rosuvastatin acid;
- c) extracting the solution with a water immiscible solvent to obtain an organic phase;
- d) evaporating the organic phase to obtain a residue;
- 10 e) combining the residue with an organic solvent and the cyclohexylamine to obtain the cyclohexylamine salt of claim 24;
- f) cooling the organic solvent to precipitate the cyclohexylammonium salt; and
- g) converting the cyclohexylammonium salt to the calcium salt.
- 15 37. The process of claim 36, wherein the mixture obtained in step a) further includes a water miscible solvent.
38. The process of claim 37, wherein the water miscible solvent is acetonitrile.
39. The process of claim 38, wherein the ratio of water to acetonitrile is about 1:2 to about 2:1 by volume.
- 20 40. The process of claim 39, wherein the ratio between the water and acetonitrile is about 1:1.
41. The process of claim 36, wherein the weight to volume ratio of the calcium salt is about 1:10 to about 1:40 Kg/L.
42. The process of claim 36, wherein the acid is selected from the group
- 25 consisting of: hydrochloride, hydrosulfuric, and hydrobromide.
43. The process of claim 36, wherein the water miscible solvent in step c) is ethyl acetate.
44. The process of claim 36, wherein the evaporation is at a pressure below about 760mmHg.
- 30 45. A process for purifying rosuvastatin calcium comprising
- a) combining rosuvastatin calcium with water to obtain a mixture;
- b) combining the mixture with an acid to obtain a solution of rosuvastatin acid;

- 5 c) extracting the solution with a water immiscible solvent to obtain an organic phase;
- d) evaporating the organic phase to obtain a residue;
- e) combining the residue with an organic solvent and an amine to obtain the ammonium salt;
- f) cooling the organic solvent to precipitate the ammonium salt; and
- g) converting the ammonium salt to the calcium salt.
46. The process of claim 45, wherein the mixture obtained in step a) further includes a water miscible solvent.
- 10 47. The process of claim 45, wherein the water miscible solvent in step a) is acetonitrile.
48. The process of claim 47, wherein the ratio of water to acetonitrile is about 1:2 to about 2:1
49. The process of claim 48, wherein the ratio of water to acetonitrile is about 1:1.
- 15 50. The process of claim 45, wherein the weight to volume ratio of the calcium salt is about 1:10 to about 1:40 Kg/L.
51. The process of claim 45, wherein the acid is selected from the group consisting of: hydrochloric hydrosulfuric and hydrobromic
52. The process of claim 45, wherein the water immiscible solvent in step c) is ethyl acetate.
- 20 53. The process of claim 45, wherein the evaporation is at a pressure below about 760mmHg.
54. The process of claim 45, wherein the organic solvent in step e) is ethylacetate.
55. The process of claim 45, wherein the amine is isopropyl amine or cyclohexylamine.
- 25 56. A process for preparing a pharmaceutical formulation of rosuvastatin calcium comprising converting the rosuvastatin calcium of claim 11, 12, 14, 33, 34, 36 or 45 to a pharmaceutical formulation.
- 30

FORM A OF ROSUVASTATIN ISOPROPYL AMMONIUM SALT

FILE: ROSU-KC429, ID:
STEP : 0.050° CNT TIME: 0.600 SEC.
RANGE: 2.00 - 40.00 (DEG) CONT. SCAN RATE : 5.00 DEG/MIN.




FIG.1

FORM B OF ROSUVASTATIN CYCLOHEXYL AMMONIUM SALT

FILE: ROSU KC-434 SHS, ID: ANALYST: LIMOR

STEP : 0.050° CNT TIME: 0.600 SEC.

RANGE: 2.00 - 40.00 (DEG) CONT. SCAN RATE : 5.00 DEG/MIN.




FIG. 2

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2004/039469

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D239/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01/60804 A (ASTRAZENECA AB; ASTRAZENECA UK LIMITED; SHIONOGI & CO., LTD; TAYLOR, N) 23 August 2001 (2001-08-23) cited in the application page 6, line 20 - line 29 claim 13	1-55
X	page 2, line 8 - line 12	56
Y	WO 00/49014 A (ASTRAZENECA UK LIMITED; SHIONOGI & CO. LTD; KOIKE, HARUO; KABAKI, MIKI) 24 August 2000 (2000-08-24) page 5, line 5 - line 8	1-55
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

15 March 2005

Date of mailing of the international search report

23/03/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kollmannsberger, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2004/039469

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with Indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>WO 00/17150-A (LEK PHARMACEUTICAL AND CHEMICAL COMPANY D.D; PFLAUM, ZLATKO) 30 March 2000 (2000-03-30) page 5, paragraph 1 page 11, paragraph 2 page 12, line 30 - page 13, line 7 examples 22,23 claim 18</p> <p>-----</p>	1-55

INTERNATIONAL SEARCH REPORT

 Int 1al Application No
 PCT/US2004/039469

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0160804 A	23-08-2001	AU 775569 B2	05-08-2004
		AU 3208401 A	27-08-2001
		BG 106969 A	30-04-2003
		BR 0108378 A	11-03-2003
		CA 2397450 A1	23-08-2001
		CN 1418198 A	14-05-2003
		CZ 20022754 A3	13-11-2002
		EE 200200445 A	15-12-2003
		EP 1263739 A1	11-12-2002
		WO 0160804 A1	23-08-2001
		HU 0204051 A2	28-05-2003
		JP 2003523334 T	05-08-2003
		NO 20023853 A	14-08-2002
		NZ 520032 A	26-03-2004
		PL 356472 A1	28-06-2004
		SK 11742002 A3	04-02-2003
		US 2003045718 A1	06-03-2003
		ZA 200205331 A	03-10-2003
WO 0049014 A	24-08-2000	AU 760145 B2	08-05-2003
		AU 2557300 A	04-09-2000
		BR 0008301 A	22-01-2002
		CA 2362594 A1	24-08-2000
		CN 1340052 A , C	13-03-2002
		CZ 20012930 A3	14-11-2001
		EE 200100430 A	16-12-2002
		EP 1155015 A1	21-11-2001
		WO 0049014 A1	24-08-2000
		HU 0200301 A2	28-08-2002
		JP 2003518474 T	10-06-2003
		NO 20013994 A	16-10-2001
		NZ 513261 A	31-10-2003
		PL 350185 A1	18-11-2002
		SK 11832001 A3	07-01-2002
		TR 200102360 T2	21-12-2001
		TR 200401874 T2	21-10-2004
		US 2004049036 A1	11-03-2004
		US 6844437 B1	18-01-2005
		ZA 200106370 A	04-11-2002
WO 0017150 A	30-03-2000	SI 20070 A	30-04-2000
		AT 271026 T	15-07-2004
		AU 765373 B2	18-09-2003
		AU 5528599 A	10-04-2000
		CA 2343646 A1	30-03-2000
		CN 1318046 A	17-10-2001
		DE 69918697 D1	19-08-2004
		EP 1466886 A2	13-10-2004
		EP 1114021 A1	11-07-2001
		HU 0103007 A2	29-05-2002
		WO 0017150 A1	30-03-2000
		IL 142055 A	20-06-2004
		JP 2002526467 T	20-08-2002
		NZ 509583 A	31-10-2003
		US 2003120086 A1	26-06-2003
		US 6583295 B1	24-06-2003
		US 2005049422 A1	03-03-2005